


Understanding Superficial Thrombophlebitis

A clot and swelling in a vein just under the skin — when it is simple, and when it needs more

A surface-vein clot is usually less dangerous than a deep clot, and it often settles with simple care. But it is not always harmless. How long the clot is and how close it sits to the deep veins decides what you need.



Online: <https://go.riasalimd.com/superficial-phlebitis-guide>

Rias KS Ali, MD FACC

Board Certified in Interventional Cardiology & Cardiovascular Disease

4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com

Last Updated: August 2026



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>

1 of 22 | Page



Save
Contact Info

Names & Terms You Will Hear

Term	Meaning
Superficial thrombophlebitis	A clot with swelling in a vein just under the skin. 'Thrombo' = clot, 'phlebitis' = vein swelling.
Superficial vein thrombosis (SVT)	The newer name doctors use. It means the same thing: a clot in a surface vein.
Phlebitis	Swelling of a vein. When a clot is also present, it is called thrombophlebitis.
Deep vein thrombosis (DVT)	A clot in a DEEP vein, down near the bone. More dangerous than a surface clot. A different problem.
Pulmonary embolism (PE)	A clot that breaks loose and travels to the lungs. A medical emergency.
Saphenofemoral junction	The spot in the groin where the main surface leg vein joins the deep vein. A clot near here is the risky kind.
Great saphenous vein	The long surface vein that runs from the inner ankle up to the groin. The most common site of a leg SVT.
Varicose veins	Bulging, twisted surface veins. Blood moves slowly in them, so clots form more easily.
Duplex ultrasound	A painless scan that shows the vein, the clot, and how close it sits to the deep system. The key test.
Cellulitis	A skin infection — red, warm, and tender like phlebitis, but caused by bacteria, not a clot. Needs antibiotics.
Fondaparinux	A once-a-day blood-thinner shot. The most-studied drug for an extensive surface-vein clot.
DOAC	A direct oral anticoagulant — a blood-thinner pill (such as rivaroxaban or apixaban). An option for some clots.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Key fact: A surface-vein clot is usually less dangerous than a deep clot, but it is **not always harmless**. A long clot, or one near where the surface vein joins the deep vein, can spread into the deep system. **How long the clot is and where it sits decide what you need — so an ultrasound is often done.**



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>




Save
Contact Info

What Is Superficial Thrombophlebitis?

- Superficial thrombophlebitis is a blood clot, plus swelling, in a vein just under the skin. The Greek words mean clot (thrombo) and vein swelling (phlebitis).
- You can usually feel it: a firm, tender, warm, red streak — like a hard cord — running along a surface vein. The skin over it may look pink or red.
- It often shows up in the leg, frequently along a varicose vein. In the arm, it often follows an IV line or a blood draw.
- It happens when blood moves too slowly, the vein wall is irritated, or the blood clots more easily. Varicose veins, IV lines, long trips, surgery, pregnancy, and some cancers all raise the chance.
- It is usually less dangerous than a deep-vein clot (DVT) and often settles with simple home care. But it is not always harmless — see why on the next pages.

Why Location Matters: Distance to the Deep Vein



The main surface leg vein runs up to meet the deep vein at the junction in the groin. A short clot low in the leg is usually simple. A long clot, or one near the junction, can reach the deep system — that is the kind that needs a blood thinner.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Why It Matters

- A surface clot is closer to the skin and farther from the lungs, so it is usually safer than a deep clot. Many cases settle in 1 to 2 weeks with simple care.
- But it is NOT always harmless. A long clot, or one in a surface vein near where that vein joins a deep vein, can grow into the deep system.
- Once a clot reaches the deep veins, it becomes a DVT — and a piece can break off and travel to the lungs (a pulmonary embolism, or PE). PE can be life-threatening.
- Because of this risk, the length of the clot and its distance from the deep-vein junction are what matter most. They decide whether you need just comfort care or a blood thinner.
- An ultrasound is often done to measure the clot, find out how close it sits to the deep veins, and make sure a silent DVT is not already present.
- Up to 1 in 4 people with a surface-vein clot also have, or go on to develop, a deep clot. That is why a careful check is worth it.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Telling them apart: phlebitis vs DVT vs cellulitis

Feature	Surface clot (phlebitis)	Deep clot (DVT)	Cellulitis (infection)
What it is	Clot in a vein under the skin	Clot in a deep vein near the bone	Bacterial skin infection
How it feels	Firm, tender, warm red cord you can feel	Whole-leg or calf swelling, deep ache	Spreading red, hot, tender skin
Danger level	Usually mild; risky if long or near junction	Can travel to the lungs (PE)	Can spread; needs antibiotics
Main test	Duplex ultrasound	Duplex ultrasound	Exam; sometimes blood tests
Treatment	Warm care, or a thinner if extensive	Full-dose blood thinner	Antibiotics



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Surface vs Deep: Why Location Matters

- Surface veins sit just under the skin. Deep veins run lower, near the bone, wrapped by muscle. A clot in each is a different problem.
- A surface clot is farther from the lungs, so it is usually safer. You can see and feel it as a red, tender cord — which is why it gets found early.
- The risk goes up when a surface clot is long, or when it sits high in the thigh near where the surface vein joins the deep vein (the saphenofemoral junction).
- Near that junction, the clot has only a short way to grow before it reaches the deep system. That is the kind that can turn into a DVT.
- So the question is not just 'is it a surface clot?' but 'how long is it, and how close is it to the deep veins?' The ultrasound answers both.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Risk Factors

Some patients are more likely to develop this condition. Knowing your risk helps your care team take extra precautions.

Risk Factor	Why It Increases Risk
Varicose veins	Blood moves slowly in stretched, twisted veins, so clots form more easily. This is the most common setting for a leg SVT.
An IV line or blood draw	A catheter or needle irritates the vein wall. This is the most common cause of an arm SVT.
Sitting still for a long time	Long flights, car trips, or bed rest let blood pool and clot. Move and stretch often.
Recent surgery or injury	Healing and bruising make blood more likely to clot, and the leg may move less.
Pregnancy and the weeks after birth	Hormones and pressure from the womb slow blood flow in the legs.
Estrogen — the pill or hormone therapy	Estrogen makes blood clot more easily. The risk is higher if you also smoke.
A past clot — SVT, DVT, or PE	Once you have had a clot, you are more likely to have another.
Some cancers and clotting disorders	Certain cancers and inherited clotting problems make clots form more easily. Repeated SVT in different spots can be a clue.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Treatment Options

- Step 1 — Check the clot. A duplex ultrasound measures how long the clot is and how close it sits to the deep-vein junction. It also rules out a hidden DVT.
- Simple cases (a short clot, low in the leg, away from the junction): warm, moist compresses, an anti-inflammatory medicine (like ibuprofen) for pain, a compression stocking, and staying active.
- Keep moving. Walking pumps blood out of the legs and lowers the chance the clot spreads. Do not lie still in bed.
- Raise the leg when resting to ease swelling. Most simple surface clots settle within 1 to 2 weeks.
- More extensive cases (a clot 5 cm or longer, or one within a few centimeters of the deep-vein junction): a short course of a blood thinner to stop the clot from spreading.
- The most-studied choice is fondaparinux, a once-a-day shot, usually for about 45 days. In the CALISTO trial it cut clot events (spread, DVT, or PE) by about 79% without raising serious bleeding.
- A DOAC pill (such as rivaroxaban) is an option for some patients instead of the shot. Your doctor weighs your bleeding risk before choosing.
- If the clot is right at the junction, or already reaches the deep vein, you are treated like a DVT — with full-dose blood thinners. Rarely, a vein near the junction is tied off in surgery.
- Treat the cause, too. If an IV line caused it, the line comes out. If varicose veins caused it, your doctor may suggest treating them later to prevent another clot.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Treatment Depends on How Long the Clot Is and Where It Sits

SIMPLE care (short, low clot)

- Warm, moist compresses on the sore area
- Anti-inflammatory pills (like ibuprofen) for pain
- Compression stocking to ease swelling
- Keep walking — do not lie still in bed

Most settle in 1-2 weeks

BLOOD THINNER (long clot or near junction)

- A short course of a thinner (fondaparinux or a DOAC)
- Usually about 45 days to stop the clot from spreading
- Used when the clot is 5 cm+ long or close to the deep vein
- Ultrasound first checks the deep veins and the junction

Stops spread to a DVT or lung clot

Treatment depends on the clot. Short, low clots get simple comfort care. Long clots (5 cm or more) or clots near the deep-vein junction get a short course of a blood thinner — usually fondaparinux for about 45 days — to stop the clot from spreading.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

When It's Simple vs When It Needs a Blood Thinner

- Simple: a short clot (under 5 cm), low in the leg, far from the deep-vein junction, with no other clot risks. These often settle with warm care, anti-inflammatory medicine, compression, and walking.
- Needs a thinner: a clot 5 cm or longer, or one within about 3 cm of the deep-vein junction. Here the spread risk is real, so a short course of a blood thinner is used.
- The usual choice is fondaparinux, a once-a-day shot, for about 45 days. In the CALISTO trial it cut clot events by about 79% versus placebo, with no rise in serious bleeding.
- A DOAC pill is an option for some patients who prefer not to use a shot. Your doctor weighs your bleeding risk and other medicines.
- If the clot is right at the junction or already reaches the deep vein, you are treated like a DVT, with full-dose blood thinners.
- See our DVT guide at go.riasalimd.com/dvt-guide and Bridging guide at go.riasalimd.com/anticoag-bridge-guide for more on blood thinners.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Comfort Measures at Home (No Medication Needed)

These simple steps support healing and ease symptoms. Use them alongside any medication your doctor prescribes.

- Put a warm, moist cloth on the sore area for 15 to 20 minutes, a few times a day. Warmth eases the ache and helps the swelling settle.
- Wear a compression stocking during the day if your doctor advises it. It supports the vein and reduces swelling.
- Walk every day. Each step pumps blood up and out of the leg and lowers the chance the clot spreads.
- Raise the leg above heart level when you rest. Use a pillow under the calf, not just the knee.
- Avoid rubbing or massaging the firm cord. Let it settle on its own.
- Mark the edge of the red area with a pen. If the redness spreads past your mark over a day, call us.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Risks, Benefits, and Alternatives

Every choice has trade-offs. Use this table to start the shared decision conversation.

Option	Risks	Benefits	Alternatives
Simple care (warm compresses, anti-inflammatory, compression, walking)	Will not stop a clot that is already spreading. Anti-inflammatory pills can upset the stomach or affect the kidneys if overused.	Eases pain and swelling fast. No bleeding risk. Right for short, low clots that are far from the deep veins.	A blood thinner if the clot is long or near the junction; ultrasound first to be sure which group you are in.
Fondaparinux (a blood-thinner shot, about 45 days)	A once-a-day shot under the skin. Small bleeding or bruising risk. Not used if kidneys are weak.	Best-studied option. Cut clot events by about 79% in the CALISTO trial. Stops spread to a DVT or lung clot.	A DOAC pill instead of the shot; simple care alone for short, low clots; full-dose thinner if it reaches the deep vein.
A DOAC pill (such as rivaroxaban)	A daily pill. Small bleeding risk. Some are not used in pregnancy or with certain other drugs.	No shots needed. Convenient. A reasonable choice for an extensive surface clot in many patients.	Fondaparinux (more trial evidence for SVT); simple care for low-risk clots.
Watchful waiting (recheck, no thinner)	A clot can quietly grow toward the deep vein. Needs a repeat ultrasound to stay safe.	Avoids any bleeding risk. Reasonable for a very short clot, low in the leg, far from the junction.	A short course of a blood thinner if a follow-up scan shows the clot is spreading.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Common Misconceptions

Myth	Reality
A surface-vein clot is always harmless — it's just a sore vein.	Usually it is mild, but not always. A long clot, or one near where the surface vein joins the deep vein, can spread into the deep system and even reach the lungs. Location and length decide the risk.
If it's not a deep clot, I don't need any tests.	An ultrasound is often worth it. Up to 1 in 4 people with a surface clot also have a hidden deep clot. The scan also measures how close the clot sits to the deep-vein junction.
Antibiotics will fix it.	A surface-vein clot is not an infection, so antibiotics do not help. Antibiotics are for cellulitis — a skin infection that can look similar. Your doctor tells the two apart.
I should rest in bed until it heals.	The opposite. Lying still lets blood pool and clot. Walking pumps blood out of the leg and lowers the chance the clot spreads. Stay active unless told otherwise.
Phlebitis and a DVT are the same thing.	No. Phlebitis is a clot in a SURFACE vein, just under the skin. A DVT is a clot in a DEEP vein near the bone. The deep clot is more dangerous and is treated differently.
Blood thinners are needed for every surface clot.	Not for short clots low in the leg, far from the deep veins — those often settle with simple care. Thinners are saved for long clots or clots near the junction, where the spread risk is real.

Related guides: The **DVT** guide at go.riasalimd.com/dvt-guide explains deep clots. The **PE** guide at go.riasalimd.com/pe-guide covers lung clots. The **Varicose Veins** guide at go.riasalimd.com/varicose-veins-guide and the **Venous Insufficiency** guide at go.riasalimd.com/vi-guide cover the leg veins where these clots often start. The **Bridging** guide at go.riasalimd.com/anticoag-bridge-guide covers blood thinners.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Possible Complications

Knowing what can go wrong helps you spot problems early. Most complications are uncommon, especially with treatment.

Where / What	What Can Happen
Spread into a deep vein (DVT)	The main worry. A surface clot can grow along the vein and reach the deep system, especially near the saphenofemoral junction. New deep swelling, calf pain, or tightness needs a same-day ultrasound.
Pulmonary embolism (PE)	If a deep clot forms and a piece breaks loose, it can travel to the lungs. Sudden shortness of breath, chest pain, or coughing up blood is an emergency — call 911.
The clot comes back	If the cause is not treated — varicose veins, a clotting problem, or an ongoing IV line — another surface clot can form. Repeated clots in different spots should prompt a closer look.
Skin staining or a firm cord that lingers	After the clot settles, the vein may stay firm for weeks and the skin over it may turn brown. This is harmless and usually fades over time.
Confusion with a skin infection	Phlebitis and cellulitis both look red and warm. Treating one as the other delays care. Spreading redness with fever points more to infection — get checked.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Warning Signs That Mean Get Checked

- Sudden shortness of breath, chest pain, a racing heart, or coughing up blood — call 911. These can mean a clot has reached the lungs (a pulmonary embolism).
- New swelling of the calf or whole leg, deep aching, warmth, or tightness — call us today or go to the ER. This can mean the clot has reached a deep vein (a DVT).
- The firm red cord is spreading UP your leg toward the groin — call us today. A clot nearing the junction needs a prompt ultrasound.
- Spreading redness with fever — call us today. This may be a skin infection (cellulitis) rather than just the clot.
- When in doubt, call. A quick ultrasound is far easier than treating a clot that has already spread. See our PE guide at go.riasalimd.com/pe-guide.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Points to Know

If you remember nothing else, remember these key points.

- Superficial thrombophlebitis is a clot, plus swelling, in a vein just under the skin. You can usually feel it as a firm, tender, warm, red cord.
- It is usually less dangerous than a deep clot and often settles with simple care: warm compresses, anti-inflammatory medicine, compression, and walking.
- But it is not always harmless. A long clot, or one near the deep-vein junction in the groin, can spread into the deep veins.
- Length and location decide your treatment. That is why an ultrasound is often done to measure the clot and check the deep veins.
- Extensive clots (5 cm or longer, or near the junction) are treated with a short course of a blood thinner — usually fondaparinux for about 45 days.
- Keep moving and raise the leg when resting. Do not rest in bed — that makes clots more likely.
- New leg swelling, calf pain, or warmth may mean a DVT — call us today. Sudden breathlessness or chest pain may mean a lung clot — call 911.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

When to Call Us - and When to Call 911

If you are not sure, call. We would rather hear from you twice than miss a real problem.

- Spreading redness, more swelling, or a fever — call us today (we need to rule out a skin infection or a spreading clot).
- New swelling, deep aching, or tightness in the calf or whole leg — call us today or go to the ER (rule out a deep clot / DVT).
- The firm cord is creeping UP toward the groin — call us today (a clot nearing the junction needs a prompt scan).
- Sudden shortness of breath, chest pain, fast heartbeat, or coughing up blood — call 911 now (possible lung clot / PE).
- Pain that is getting worse despite warm compresses and anti-inflammatory medicine — call us to be rechecked.
- You are on a blood thinner and notice unusual bruising, bleeding gums, or blood in urine or stool — call us today.
- A red, tender vein appears where an IV line was placed — tell your care team so the line can be removed and the area watched.

Office: (727) 943-5200

Call 911 now for sudden shortness of breath, chest pain, a racing heart, or coughing up blood — a clot may have traveled to the lungs (a pulmonary embolism). **Call us today** for new calf or whole-leg swelling, deep aching, or warmth — this can mean a deep clot (DVT). When in doubt, get checked: a quick scan beats a missed clot.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Trusted Resources

The links below are the trusted resources Dr. Ali used to write this guide and that he recommends you visit for more information. Each is a clickable link in the digital version and a Short.io tracked URL on the printed version.

- [Cleveland Clinic — Thrombophlebitis](#) - Plain-language overview of a surface-vein clot, its symptoms, simple care, and when it points to a deeper clot.
- [Mayo Clinic — Thrombophlebitis](#) - Symptoms, causes, and risk factors, plus how surface phlebitis relates to DVT.
- [MedlinePlus — Superficial Thrombophlebitis](#) - NIH/NLM patient overview of causes, signs, and treatment.
- [American Heart Association — Venous Thromboembolism](#) - How vein clots form and why a deep clot is more dangerous than a surface one.
- [AAFP — Superficial Thrombophlebitis](#) - A primary-care guide to telling phlebitis apart from DVT and cellulitis.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Sources Used to Build This Guide

Every recommendation in this guide is grounded in one or more of the studies, guidelines, or trusted resources below. Short links are tracked so Dr. Ali can see which references patients are reading; nothing personally identifying is captured.

- [Cleveland Clinic — Thrombophlebitis](#) [Patient Education]
Plain-language overview of a clot and inflammation in a surface vein, symptoms, warm-compress care, and when it signals deeper clot risk.
- [Mayo Clinic — Thrombophlebitis](#) [Patient Education]
Patient-facing symptoms, causes, risk factors, and the relationship between superficial phlebitis and DVT.
- [MedlinePlus — Superficial Thrombophlebitis](#) [Patient Education]
NIH/NLM overview of causes, presentation, and treatment to anchor the explainer in plain English.
- [CHEST Antithrombotic Therapy for VTE Disease — CHEST Guideline \(Stevens 2021\)](#) [Guideline]
Authoritative basis for when anticoagulation is indicated for superficial vein thrombosis (length, proximity to the deep system) versus supportive care.
- [ASH 2020 Guidelines for VTE: Treatment of DVT and PE \(Blood Advances\)](#) [Guideline]
American Society of Hematology recommendations: whole-leg ultrasound, fondaparinux 2.5 mg for SVT 5 cm or longer and 3 cm or more from the junction, and therapeutic anticoagulation when the clot is within 3 cm of the deep-vein junction.
- [CALISTO Trial — Fondaparinux for Superficial-Vein Thrombosis \(NEJM 2010\)](#) [Clinical Trial]
Landmark randomized trial: 45 days of fondaparinux 2.5 mg cut symptomatic clot events (DVT, PE, or clot extension) by about 79% versus placebo without raising major bleeding. The evidence behind treating extensive SVT with a blood thinner.
- [CALISTO Trial — PubMed Record \(Decousus 2010\)](#) [Clinical Trial]
PubMed citation for the CALISTO study (3,002 patients) used to verify the 45-day course, 2.5 mg dose, and event-reduction figure.
- [American Heart Association — Venous Thromboembolism](#) [Patient Education]
AHA venous-clot overview used to frame how a surface-vein clot relates to the deeper, more dangerous DVT and PE.
- [AAFP — Superficial Thrombophlebitis Diagnosis and Treatment \(American Family Physician\)](#) [Review]
Primary-care review of how to tell SVT apart from cellulitis and DVT, the role of duplex ultrasound, and supportive-care versus anticoagulation decisions.
- [StatPearls — Superficial Thrombophlebitis \(NCBI Bookshelf\)](#) [Reference]
Peer-reviewed reference on causes (varicose veins, IV catheters), the saphenofemoral junction risk, and management thresholds.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

About This Guide

Rias KS Ali, MD FACC | Board Certified in Interventional Cardiology & Cardiovascular Disease

4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com

Version: 2026-08-15

Reviewer note: Reviewed against Cleveland Clinic, Mayo Clinic, and AAFP patient resources, with the ASH 2020 and CHEST VTE guidelines and the CALISTO fondaparinux trial. This guide adds: a surface-vs-deep depth diagram, a saphenofemoral-junction map showing why location matters, a treatment-by-extent panel, a side-by-side table telling phlebitis apart from DVT and cellulitis, and cross-links to companion venous guides (DVT, PE, Varicose Veins, Venous Insufficiency, Bridging).

Disclaimer

This guide is for educational purposes only and does not replace medical advice from your physician. Recommendations may not apply to every patient. Always follow the specific instructions of Dr. Ali and your care team. In an emergency call 911 or go to the nearest emergency department.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info

Take This Guide With You



Scan or visit: <https://go.riasalimd.com/superficial-phlebitis-guide>

Online: <https://go.riasalimd.com/superficial-phlebitis-guide>

Office: (727) 943-5200

4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/superficial-phlebitis-guide>



Save
Contact Info