


Understanding Xarelto (Rivaroxaban)

A blood thinner used for stroke prevention, clot treatment, and heart disease

Stops clots before they start. Take with food at the right dose. Never stop without calling US.



Online: <https://go.riasalimd.com/xarelto-guide>

Rias KS Ali, MD FACC

Board Certified in Interventional Cardiology & Cardiovascular Disease

4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com

Last Updated: August 2026



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Names & Terms You Will Hear

Term	Meaning
Xarelto	The brand name for rivaroxaban in the United States.
Rivaroxaban	The generic name. Available as 2.5 mg, 10 mg, 15 mg, and 20 mg tablets.
Direct oral anticoagulant (DOAC)	A newer class of blood thinners that work directly on one clotting factor. Unlike warfarin, DOACs do not require regular INR blood tests.
Factor Xa inhibitor	The mechanism — rivaroxaban blocks Factor Xa, a key step in the clotting cascade, preventing clots from forming.
Anticoagulant / blood thinner	A medicine that makes it harder for blood to clot. It does NOT dissolve existing clots — it prevents new ones.
INR (International Normalized Ratio)	The blood test used to monitor warfarin. Rivaroxaban does NOT require INR monitoring.
DVT / PE	Deep vein thrombosis (clot in a leg vein) and pulmonary embolism (clot in the lung) — two of the main conditions rivaroxaban treats or prevents.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

What Is Xarelto (Rivaroxaban)?

- Rivaroxaban (Xarelto) is a blood thinner that blocks Factor Xa — a key step in the clotting cascade — preventing dangerous clots from forming.
- Unlike warfarin, it does not require regular INR blood tests. The dose is fixed based on your condition, not a weekly blood draw.
- It comes in four strengths: 2.5 mg (CAD/PAD vascular dose), 10 mg (post-surgery or extended VTE prevention), 15 mg (DVT/PE treatment first 3 weeks), and 20 mg (AFib and VTE treatment).
- Rivaroxaban does NOT dissolve clots that are already there — it prevents new clots from forming while your body resolves the existing clot.
- If serious bleeding occurs, there is a reversal agent: andexanet alfa (Andexxa). Emergency departments stock it. Carry a medical ID card stating you take rivaroxaban.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Why It Matters

- In the ROCKET AF trial (14,000 patients with AFib), rivaroxaban was as effective as warfarin for stroke prevention with less intracranial bleeding.
- In the EINSTEIN trials, rivaroxaban was as good as the LMWH-to-warfarin regimen for treating DVT and PE — with the convenience of one oral drug from day one.
- In COMPASS, the 2.5 mg twice-daily vascular dose (added to low-dose aspirin) cut heart attack, stroke, and cardiovascular death by 24% in stable CAD/PAD patients.
- Blood clots in AFib are the leading cause of large, disabling strokes. Without anticoagulation, AFib patients have a 3-5x higher stroke risk.
- Stopping rivaroxaban without a plan creates a clot-rebound window — the highest-risk period for stroke or PE is the first 2-4 weeks after stopping.




Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info



A real deep vein thrombosis (DVT) — unilateral leg swelling and redness from a clot in the deep veins. This is one of the conditions the EINSTEIN trials tested rivaroxaban against: 15 mg twice daily for the first 21 days, then 20 mg once daily, to treat the clot and prevent it from traveling to the lungs (pulmonary embolism). Image: James Heilman, MD, Wikimedia Commons (CC BY-SA 3.0).



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Treatment Options

- AFib stroke prevention: 20 mg once daily with your evening meal (food increases absorption significantly).
- DVT or PE treatment: 15 mg twice daily with food for the first 21 days, then 20 mg once daily with food for the remainder of treatment.
- Post-hip or knee replacement: 10 mg once daily, starting 6-10 hours after surgery, for 35 days (hip) or 12 days (knee).
- CAD/PAD vascular dose: 2.5 mg twice daily plus low-dose aspirin (100 mg once daily) — this is a lower-than-anticoagulation dose, used only for the vascular indication.
- Never stop rivaroxaban without calling us first — stroke or clot recurrence risk peaks in the first weeks after stopping.
- If you need surgery or a dental procedure, call us at least 48-72 hours in advance to discuss a planned hold and restart timing.
- Bleeding risk is real but manageable: use a soft toothbrush, electric razor, and avoid NSAIDs (ibuprofen, naproxen) unless we specifically approve them.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Risks, Benefits, and Alternatives

Every choice has trade-offs. Use this table to start the shared decision conversation.

Option	Risks	Benefits	Alternatives
Rivaroxaban (Xarelto)	Bleeding — anywhere, including GI or intracranial (rare). No routine INR monitoring but no easy dose adjustment. Reversal agent exists but is expensive. Drug interactions with strong CYP3A4/P-gp drugs.	No INR testing. Once-daily for most indications. Effective as warfarin for AFib/VTE. COMPASS vascular-dose benefit in CAD/PAD. Multiple approved indications.	Apixaban (Eliquis — often less GI bleeding), dabigatran (Pradaxa), warfarin (INR monitoring required), no anticoagulation (high stroke/clot risk).
Apixaban (Eliquis)	Bleeding. Twice-daily dosing. No INR required. Reversal agent (andexanet alfa) available.	Meta-analyses show slightly less GI bleeding vs rivaroxaban. Twice-daily dosing may improve steady-state levels.	Rivaroxaban (once daily for most indications), warfarin, dabigatran.
Warfarin (Coumadin)	Bleeding. Weekly INR tests. Highly food- and drug-interactive. Narrow therapeutic window — hard to stay in range.	Decades of data. Cheap. Two reversal options (vitamin K, FFP/4F-PCC). Required for mechanical heart valves — DOACs cannot be used.	DOACs (preferred over warfarin in nonvalvular AFib and VTE per guidelines). Rivaroxaban or apixaban preferred for most patients.
No anticoagulation in AFib	3-5x higher annual stroke risk. Strokes from AFib clots are typically large and disabling.	No drug side effects or bleeding risk from medication.	Anticoagulation is Class I (highest) recommendation in AFib with CHA ₂ DS ₂ -VASc score of 2 or higher (men) or 3 or higher (women).



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Common Misconceptions

Myth	Reality
I can stop Xarelto if I feel fine and my heart rhythm is normal.	Stopping on your own is dangerous. Even when AFib is not felt, it can be running silently — and stopping anticoagulation creates a rebound period of high clot risk. Always call us before stopping.
I need weekly INR tests like my neighbor who takes warfarin.	Rivaroxaban does NOT require INR monitoring. The dose is fixed. Warfarin levels fluctuate with diet and other drugs — rivaroxaban does not. This is one of its main advantages.
Xarelto thins my blood — my blood is watery.	The blood itself is unchanged. Rivaroxaban slows one step in the clotting cascade. Your blood cells, platelets, and vessels are all normal. You may bruise more easily or bleed longer from a cut, but your blood is not thinner in a physical sense.
Missing one dose is not a big deal.	For AFib, missing doses increases stroke risk. If you realize on the same day, take it. If it's almost time for the next dose, skip. Never double-dose. Set a phone alarm — consistency matters more than the exact time.
If I take aspirin, I don't need Xarelto.	Aspirin and blood thinners work differently. For AFib, aspirin does NOT reliably prevent clot-based strokes. The standard of care is a DOAC like rivaroxaban, not aspirin. For the COMPASS CAD/PAD indication, both are prescribed together at low doses — that is a special combined regimen.
I should avoid all vegetables and greens because of my blood thinner.	That restriction applies to warfarin, not rivaroxaban. Rivaroxaban is not affected by vitamin K in food. Eat a normal, balanced diet. Grapefruit should still be limited as it affects metabolism.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Myth	Reality
There is no antidote if I have a major bleed.	Andexanet alfa (Andexxa) is an FDA-approved reversal agent for rivaroxaban. Most major hospitals stock it. You should also carry a medical ID card so that ER staff know immediately what you take.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
 4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
 Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Possible Complications

Knowing what can go wrong helps you spot problems early. Most complications are uncommon, especially with treatment.

Where / What	What Can Happen
Major bleeding	The most serious risk. Risk is ~2-3% per year for major bleeding events. Intracranial hemorrhage occurs in <0.5% per year — lower than with warfarin. GI bleeding is slightly higher than warfarin but lower than dabigatran.
Surgical / procedural bleeding	Rivaroxaban must be held before most invasive procedures (24-48 hours for elective; longer for high-risk procedures). Failure to coordinate with your care team is a preventable risk.
Drug interactions — CYP3A4 and P-gp pathway	Rivaroxaban depends on both CYP3A4 and P-glycoprotein for metabolism and excretion. Strong inhibitors (azole antifungals, HIV protease inhibitors) raise levels. Strong inducers (rifampin, carbamazepine, St. John's wort) lower levels unpredictably.
Rebound thrombosis on abrupt discontinuation	Stopping rivaroxaban suddenly in AFib patients creates a 2-4 week window of elevated stroke risk. Always taper or bridge under physician guidance.
Spinal hematoma	Rare but serious risk with neuraxial anesthesia (spinal or epidural blocks). Anesthesiologists must know you are on rivaroxaban before any regional anesthesia procedure.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Points to Know

If you remember nothing else, remember these key points.

- 20 mg and 15 mg tablets must be taken with food — absorption drops by 30% without a meal.
- Carry a wallet card or wear a medical alert bracelet stating you take rivaroxaban.
- Call us at least 48-72 hours before any planned surgery, procedure, or dental extraction.
- Avoid NSAIDs (ibuprofen, naproxen, Aleve, Advil) unless specifically approved by Dr. Ali — they significantly raise bleeding risk.
- Grapefruit and grapefruit juice may increase rivaroxaban levels — limit or avoid.
- Store at room temperature, away from moisture. Keep in the original bottle.
- If you need to crush a tablet (15 mg or 20 mg), mix with 50 mL of water and take with food immediately.
- Tell every provider — surgeon, dentist, any urgent care — that you are on rivaroxaban before any procedure.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

When to Call Us - and When to Call 911

If you are not sure, call. We would rather hear from you twice than miss a real problem.

- Unusual or heavy bleeding that does not stop — call 911 if severe.
- Blood in urine (pink, red, or brown) — call us today.
- Black or tarry stools, or vomiting blood — call 911 or go to the ER.
- Coughing up blood — call 911.
- Sudden severe headache, vision changes, weakness on one side, or trouble speaking — call 911 (stroke symptoms).
- Sudden chest pain or shortness of breath — call 911.
- Any fall with head injury — go to the ER and tell them you take rivaroxaban.
- Before any surgery, procedure, or if you want to stop for any reason — call us first.

Office: (727) 943-5200



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Trusted Resources

The links below are the trusted resources Dr. Ali used to write this guide and that he recommends you visit for more information. Each is a clickable link in the digital version and a Short.io tracked URL on the printed version.

- [MedlinePlus — Rivaroxaban](#) - NIH-curated patient drug information.
- [Cleveland Clinic — Rivaroxaban \(Xarelto\)](#) - Patient-friendly drug profile and safety guide.
- [American Heart Association — Anticoagulation](#) - AHA patient page on blood thinners and AFib.
- [National Blood Clot Alliance](#) - Patient advocacy site for DVT/PE education and peer support.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Sources Used to Build This Guide

Every recommendation in this guide is grounded in one or more of the studies, guidelines, or trusted resources below. Short links are tracked so Dr. Ali can see which references patients are reading; nothing personally identifying is captured.

- [ROCKET AF — Rivaroxaban vs Warfarin in Nonvalvular AFib \(Patel et al, NEJM 2011\)](#) [Clinical Trial]
Pivotal RCT: rivaroxaban noninferior to warfarin for stroke prevention in AFib; less intracranial hemorrhage. Foundation for AFib indication.
- [EINSTEIN DVT/PE — Rivaroxaban for DVT and PE \(Bauersachs et al / Büller et al, NEJM 2010/2012\)](#) [Clinical Trial]
Landmark trials establishing single-drug rivaroxaban for DVT and PE treatment, replacing the LMWH-to-warfarin bridging regimen.
- [COMPASS Trial — Rivaroxaban + Aspirin in CAD/PAD \(Eikelboom et al, NEJM 2017\)](#) [Clinical Trial]
Vascular dose (2.5 mg BID + aspirin) reduced MACE by 24% and death by 18% in stable CAD/PAD. Anchors the CAD/PAD indication.
- [RECORD Trials — Rivaroxaban after hip/knee replacement \(Eriksson et al, NEJM 2008\)](#) [Clinical Trial]
Demonstrated superiority over enoxaparin for VTE prevention post-orthopedic surgery. Foundation for the surgical prophylaxis indication.
- [ACC/AHA/HRS AFib Guideline \(January et al, Circulation 2019\)](#) [Guideline]
Class I recommendation for DOACs (including rivaroxaban) over warfarin for nonvalvular AFib. Positions rivaroxaban in the treatment algorithm.
- [ACCP VTE Guidelines — Antithrombotic Therapy for VTE Disease \(Chest 2021\)](#) [Guideline]
Strong recommendation for DOACs over vitamin K antagonists for treatment of DVT/PE. Positions rivaroxaban as a first-line option.
- [MedlinePlus — Rivaroxaban](#) [Clinical]
NIH-curated patient drug information including dosing, storage, and interactions.
- [Cleveland Clinic — Rivaroxaban \(Xarelto\)](#) [Clinical]
Patient-friendly drug profile with dosing guidance and bleeding warning signs.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

About This Guide

Rias KS Ali, MD FACC | Board Certified in Interventional Cardiology & Cardiovascular Disease

4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com

Version: 2026-08-15

Reviewer note: Reviewed against Xarelto manufacturer PI, MedlinePlus, and Cleveland Clinic drug page. Ours adds: dose-by-indication visual reference table, explicit comparison to warfarin (no INR, but no reversal button), the COMPASS vascular-dose distinction (2.5 mg BID + aspirin vs full anticoagulation doses), and the missed-dose protocol for twice-daily regimens.

Disclaimer

This guide is for educational purposes only and does not replace medical advice from your physician. Recommendations may not apply to every patient. Always follow the specific instructions of Dr. Ali and your care team. In an emergency call 911 or go to the nearest emergency department.



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info

Take This Guide With You



Scan or visit: <https://go.riasalimd.com/xarelto-guide>

Online: <https://go.riasalimd.com/xarelto-guide>

Office: (727) 943-5200

4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com



Visit
RiasAliMD.com

This guide is educational only. Always follow your doctor's specific recommendations.
4740 Mile Stretch Drive, Holiday FL 34690 | Office: (727) 943-5200 | Fax: (727) 943-5201 | RiasAliMD.com
Online: <https://go.riasalimd.com/xarelto-guide>



Save
Contact Info